

Set - A

1. Number of moles of solute dissolved in 1000 g of water is
a) molality b) molarity c) mole fraction d) % (w/v)
2. Enthalpy of a compound is equal to its
a) Heat of combustion b) Heat of formation
c) Heat of solution d) Heat of dilution
3. The atomic number is changed by which type of radioactive decay?
a) β b) α c) γ d) All of the above
4. Which of the following compound is known as Schweizer's reagent?
a) $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ b) CaSO_4
c) $[\text{Cu}(\text{NH}_3)_4]\text{SO}_4$ d) Anhydrous Copper
5. Which of the following is not true regarding crystal field theory?
a) Action betⁿ metal & ligands is electrostatic
b) Ligand are treated as point charge
c) There is no orbital interaction betⁿ ligand & metal.
d) Hybridization explain the shape of the complexes.
6. Ether is always purified before distillation because,
a) It is highly poisonous in nature.
b) It forms poisonous phosgene gas
c) It is converted into explosive peroxide
d) All of above.

7. Which of the following is an organometallic compound?
a) CH_3ONa b) CH_3SNa c) CH_3MgCl d) All of above
8. Solubility of a salt M_2X_3 is " x " mole l^{-1} . The solubility product of the salt will be
a) x^5 b) $16x^2$ c) $96x^5$ d) $108x^5$
9. Alcoholysis of acid anhydride gives
a) carboxylic acid & ester b) Ester & alcohol
c) carboxylic acid & alcohol d) only ester
10. An organic compound 'A' reacts with nitrous acid to form N-methyl-N-nitrosmethanamine. A can be obtained by the reduction of
a) Propanenitrile b) Methylisocyanate
c) Ethylisocyanide d) Propylisocyanide.
11. Given, $E^\circ_{\text{Cr}/\text{Cr}^{+3}} = 0.74\text{V}$, $E^\circ_{\text{Fe}^{+2}/\text{Fe}} = 0.42\text{V}$. The standard cell potential for the cell $\text{Cr}/\text{Cr}^{+3} (0.1\text{M}) // \text{Fe}^{+2} (0.01\text{M}) / \text{Fe}$ is,
a) 1.14V b) 0.492V
c) 0.329V d) -0.26V

b) could we use sodium hydroxide to separate zinc and silver from each other? If yes how? (1)

⇒

c) what happens when, zinc is exposed to moist air for long time? (1)


⇒ The action of air on zinc metal forms a very thin coating of zinc oxide all over it so it becomes dull.

2 a) Define polymerization.

⇒ Polyme The process of formation of polymers from monomers is known as polymerization.

b) Write the reaction involved in the preparation of polystyrene. (2)

⇒



c) what is an ideal requirement for a dye? (1)

⇒ It must have suitable attractive colour (ie.) it should absorb light in the visible region.

d) Azobenzene is a colored compound but not a dye. Explain. (1)

⇒ because it is unable to fix itself to a fibre.

3 Transition elements are defined as elements that have partially filled d-orbitals.

a) What is d-orbital degeneracy? (1)

⇒ Degenerate orbitals are the orbitals of the same subshell having equal energies. When the electrons of the orbitals are not influenced by any external factors like an electric field or magnetic field, they have same energies.

b) Give a possible reason for a fact that transition metal have high heat of atomization. (1)

⇒ Transition elements have high effective nuclear charge and a large number of valence electrons. Therefore, they form very strong metallic bonds which result in high enthalpies of atomization.

c) A transition metal forms alloy with other transition metal easily, Explain (1).

⇒ Transition metals have very similar atomic sizes. One metal can easily replace the other metal from its lattice to form solid solution (alloy). Transition metals are miscible with one another in the molten state. The molten state solution of two or more transition metals on cooling forms alloy.

d) $K_4 [Fe (CN)_6]$, potassium hexacyanoferrate(II) is a complex salt formed by iron. Write the complex ion present in it. (1)

⇒ oxidation no. of K is +1, CN = -1, Fe = +2
oxidation no. of the 4K = $+1 \times 4 = +4$, oxidation no. of the 6CN = $-1 \times 6 = -6$, oxidation no. of the Fe = +2
So, the charge of the complex ion of potassium fer.


e) Why $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$ is violet in color in $[\text{Ti}(\text{H}_2\text{O})_6]\text{Cl}_3$, but when water molecules are removed it becomes colorless? (1)

⇒


When light corresponding to the energy of the yellow-green region is absorbed by the complex, this would excite the electron from t_{2g} level to the e_g level ($t_{2g}^1 e_g^0 \rightarrow t_{2g}^0 e_g^1$). Thus the complex is violet in color.

4 The sum of the powers of concentration terms in the rate law equation is called the order of reaction.

a) Write the three possible rate laws for the given second order reaction.



→ The given second order reaction is



The possible three rate laws are listed below:

$$\text{Rate} = k[A]^1[B]^1 \quad \therefore \text{order} = 1+1 = 2$$

$$\text{Rate} = k[A]^2[B]^0 \quad \therefore \text{order} = 2+0 = 2$$

$$\text{Rate} = k[A]^0[B]^2 \quad \therefore \text{order} = 0+2 = 2$$

5) solution,

$$\text{Rate } R = k[P]' [Q]^2$$

i) $\text{Rate} = k[P]' [3Q]^2$

$$= 9R$$

∴ 9 times of initial rate

ii) $R' = Q' = 2$ (double the initial rate)

$$R' = k [2P]' [2Q]^2$$

$$= 8R$$

= 8 times of initial rate.

5 cement is one of the major component of construction field. Nepal has many cement factories. Limestone is a dominant raw material used for the manufacture of cement. When cement comes in contact with water, it sets to hard mass showing exothermic reaction.

a) You are asked to bring 1500 tons of cement for the dam construction of 400 MW hydropower station. What type of cement do you prefer and why? (2)

⇒ I will prefer PPC type of cement because PPC is cheaper & has low initial setting strength compared to OPC but hardens over a period with proper curing. It has a slow hydration process and thus generates less heat than OPC. There it is suitable for mass ~~concrete~~ ^{concrete} casting and makes the concrete more durable.

b) While manufacturing the Portland cement, the % composition of MgO is not exceeded than 6%, why?

⇒ ~~Due to~~ cement should be expensive & due to cracking

c) After preparing the cement clinker it is cooled to a temperature of 60-150°C before grinding, why? (1)

⇒ Because in the clinker cooler, the hot clinker is rapidly cooled by air in order to obtain a high degree of efficiency.


d) What is the role of Fe₂O₃ in the portland cement? (1)

⇒ To increase the hardened cement stone flexural strength, and slightly decrease the compressive strength.


To impart characteristic grey colour, strength & hardness.

6 In an



a) Perkin condensation





b) Hell-Volhard-Zelinsky reaction




c) Wolff-Kishner reduction



d) Clemmenson's reduction




e) Cannizzaro's reaction



QTR


Give a suitable reaction for the prepⁿ of ethanoic acid from

a) Ethanenitrile





b) Methyl magnesium chloride
 \Rightarrow

c) Ethanol



Also, convert ethanoic acid to propanoic acid



The compound (A) gives H_2 gas with sodium metal, yellow ppt with alkaline iodine is give aldehyde having molecular formula $\text{C}_2\text{H}_4\text{O}$ Oxidation with $\text{K}_2\text{Cr}_2\text{O}_7$

Date _____
Page _____

Q8 12 gm of impure Zn is made to react with excess of dilute H_2SO_4 . The total volume of H_2 gas liberated was found to be 4.2 L at 570 mm of Hg pressure & 273 K temperature. Determine the percentage purity of zinc.

⇒ Solution

Here, volume of H_2 liberated at NTP

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$\text{or, } \frac{760 \times V_1}{273} = \frac{570 \times 4.2}{273}$$

$$\therefore V_1 = 3.08 \text{ L}$$

When Zn react with dil. H_2SO_4 then,

$$\underset{\text{1 mole}}{Zn} + H_2SO_4 \rightarrow ZnSO_4 + \underset{\text{1 mole}}{H_2}$$

Here,

1 mole H_2 is liberated from 1 mole Zn i.e.,
22.4 L of H_2 = 65 gm of zinc
4 L of H_2 = $\frac{65}{22.4} \times 3.08$ gm of zinc
= 8.9 gram of zinc

Therefore,


$$\% \text{ purity} = \frac{\text{Pure weight}}{\text{given weight}} \times 100\%$$

$$= \frac{8.9}{12} \times 100\%$$


$$= 74.16\% \text{ ,,}$$

G-C


g) Write the principle reactions of oxidation, chlorination & hydrolysis process using ethyl alcohol.



Chlorination




Hydrolysis



b) Why is chloroform always stored in a dark colored bottle?


=> Chloroform is slowly oxidised by air in the presence of light to an extremely poisonous gas, carbonyl chloride, also known as phosgene. It is therefore stored in closed dark coloured bottles completely filled so that air is kept out.



c) It is used as an anaesthetic in past but now discouraged. why?


=> chloroform is not used as an anaesthetic these days because it may contain some poisonous phosgene due to oxidation by sunlight.

d) chloroform forms chloroacetone drug with acetone in presence of aqueous alkali. write the chemical reaction & mention any one use of this drug.




The use of chloroacetone is it acts as a hypnotic and nervous sedative.

e) what happens when, chloroform reacts with conc. Nitric acid? write an important application of such product.



It is used in agriculture as a soil fumigant. It was used in large quantities during world war I & was stockpiled during world war II.




(insecticide & war gas)



Remaining

7
=>

Hence, the compound (A) must be alcohol containing two carbon atoms i.e. ethanol ($\text{CH}_3\text{CH}_2\text{OH}$)



Hence, the organic compound (A) is ethanol.

10 a) The solubility product of barium sulphate ($BaSO_4$) is 1×10^{-10} at 298K. calculate its solubility in

- i) Pure water
- ii) $1 \times 10^{-3} M$ ~~S~~ H_2SO_4 solution.

i) \Rightarrow $BaSO_4$ ionized as,
 $BaSO_4 \rightleftharpoons Ba^{++} + SO_4^{--}$
 Here, $\alpha = 1, \beta = 1, K_{sp} = 1 \times 10^{-10}$
 $S = ?$


Now we have,


$$S = \left(\frac{K_{sp}}{\alpha^\alpha \beta^\beta} \right)^{\frac{1}{\alpha+\beta}}$$

$$= \left(\frac{1 \times 10^{-10}}{1^1 \cdot 1^1} \right)^{\frac{1}{2}}$$

$$= 1 \times 10^{-5} \text{ mole } L^{-1}$$

ii) $1 \times 10^{-3} M$ ~~S~~ H_2SO_4 solution,





b) write the eqⁿ for the formation of naphthalene
calculate the increase in entropy in the
evaporation of 1 mole of water at 100°C.
[Latent heat of vaporization of water is 2.26 kJ]

⇒ The eqⁿ for the formation of naphthalene is $C_{10}H_8$.

For entropy change of vaporization can be
given as,
$$\Delta S_v = \frac{\Delta H_v}{T}$$

Given values are, $\Delta H_v = 2.26 \text{ kJ/gm}$
 $= 2.26 \times 10^3 \text{ J/kmol}$


$$T = 373 \text{ K}$$

$$\text{Therefore } \Delta S_v = \left(\frac{2.26 \times 10^3}{373} \right)$$

$$\therefore \Delta S_v = 8.273 \text{ J/kmol}$$

44

a) Give the preparation of aniline from nitrobenzene.



b) Aniline is less basic than aliphatic amine.?

=> In case of aniline due to conjugation the lone pair density is less than that of methylamine. Due to this reason, aniline is less basic than aliphatic amine.

c) Why is it necessary to protect -NH₂ group before nitration of aniline?

=> The hurdles for the nitration of aniline are overcome by the protection of amino group by acetylation. The acetyl group reduces the reactivity of the ring & thus its oxidation does not occur easily with nitric acid HNO₃.

d) What is diazotization reaction? Why is diazotization always carried out at ice cold temperature?


=> Diazotization reaction involves the formation of diazonium salts when aromatic amines are made to react with nitrous acid in presence of mineral acid.

Diazotization is usually carried out at low temperature betⁿ 0-5°C. We have to maintain low temperature because if the temperature is above 5°C, diazonium salts which are in aqueous solution tend to decompose explosively.


e) Convert, Aniline into

i) p-aminodiazobenzene (ii) Acetanilide.

i)



ii)



Set - B

1 The unit of rate constant for zero order reaction is

- a) $\text{mol}^{-1}\text{Ls}^{-1}$ b) mol s^{-1} c) $\text{mol L}^{-1}\text{s}^{-1}$ d) L s^{-1}

2 When an alkyl halide reacts with sodium alkoxide to form symmetrical as well as unsymmetrical ethers. This reaction is called

- a) Hoffmann's reaction b) Reimer-Tiemann's reaction
c) Kolbe's reaction d) Williamson's reaction.

3 The curve obtained by plotting the pH of the solution during titration against the volume of alkali added from burette is known as a titration curve.

The curve ABCGH in the following curve represents

- a) Titration curve of strong acid vs strong base
- b) Titration curve strong acid vs weak base
- c) Titration curve of weak acid vs strong base
- d) Titration curve of weak acid vs weak base

4 An adiabatic process is the one in which

- a) system is closed to energy transfer
- b) system is closed to heat transfer
- c) The system is closed to both heat & energy transfer
- d) There is no enthalpy change.

5 Haloarenes are less reactive than benzene toward electrophilic substitution reaction due to

- a) positive inductive effect
- b) Resonance effect
- c) Negative inductive effect
- d) steric effect.

6 Which of the following is generally used to prepare paper?

- a) Protein b) Fibers c) cellulose d) vitamins.

7 All transition element exhibit in general electronic configuration of

- a) $(n-1)d^{1-10} ns^{1-2}$ b) $(n-1)d^0 ns^{1-2}$
- c) $ns^2 (n-1)d^{10}$ d) none of above

8. Which of the following is the normality of 50ml of HCl required to neutralize 10gm of CaCO_3 ?
a) 2N b) 4N c) 5N d) 1N
9. Based on the systematic formula of Iron carbonyl, $\text{Fe}(\text{CO})_5$, its conductivity is expected to be - - -
a) zero b) one c) five d) none
10. Which of the following reagent is used to make nitroethane from haloethane?
a) Alc. AgNO_3 b) Alc. AgNO_2 c) Alc. KNO_2 d) Alc. NaNO_2
11. An acid chloride (A) on treating with methyl magnesium chloride followed by acid hydrolysis gives 2-methyl propan-2-ol as a major product. Which of the following is acid chloride (A)?
a) methylanoyl chloride b) propanoyl chloride
c) ethanoyl chloride d) None of them.

G-B

1. Define the following terms:
a) Isolated system \Rightarrow A system that does not allow the exchange of either energy or matter with the surroundings is called isolated system.

b) Internal energy \Rightarrow The total sum of different form of energy associated with a particle in a system is known as Internal energy. It is denoted by a symbol ' E ' or ' U '.

c) Enthalpy \Rightarrow Enthalpy is the heat content of the system at constant pressure. It is denoted by a symbol ' H '.

d) Entropy \Rightarrow Entropy is the measurement of randomness or disorder of the molecules present in the system. It is denoted by a symbol ' S '.

e) Isochoric process \Rightarrow When a thermodynamic process proceed from one state to another state at a constant volume then it is known as Isochoric process.

Q: Sulphide ore is used in the extraction of mercury. Aristotle had named this element liquid silver.



a) write the chief ore of mercury.

\Rightarrow cinnabar (HgS)

b) why does mercury forms alloy with gold & silver?

\Rightarrow

c) How do you convert mercury into Nessler's reagent?
 ⇒ Alkaline solution of potassium tetra iodo mercurate (II), i.e. solution of K_2HgI_4 in KOH or NaOH is called Nessler's reagent. When mercuric chloride reacts with a potassium iodide solution, a yellow or scarlet red precipitate of HgI_2 is formed first which dissolves in excess KI to form soluble complex potassium tetraiodo mercurate (II)





d) Though mercury is metal, it exists in liquid state at room temperature. Why?

⇒ Mercury has a unique electron configuration which strongly resists removal of an electron, making it behave similarly to noble gas elements. As a result, mercury forms weak bonds & is a liquid at room temperature.

e) Instead of electrolytic refining, mercury is purified by treating it with 5% dil. HNO_3 . Write reaction involved in it. ~~++++~~

⇒




Hence, at anode oxygen is formed

3) a) Differentiate between

Condensation	Addition Polymers
i) It involves two different monomers.	i) It involves one monomer.
ii) Condensation of monomers results in polymers.	ii) Addition of monomers results in polymers.
iii) Common examples of condensation polymerization are nylon, silicon, etc.	iii) Common examples of addition polymers are PVC, polyethylene, Teflon, etc.

b) Why is nylon-6,6 called a copolymer? (1)
 ⇒ Because it is obtained by the condensation of hexamethylene diamine & adipic acid. Both monomers contain 6 carbons so it is called Nylon-6,6.

c) Which polymer is obtained when phenol is treated with formaldehyde in a basic medium? Show the pertinent reaction.
 ⇒ Bakelite.




4

$$E^{\circ}_{Zn/Zn^{++}} = 0.76 \text{ V}$$

$$E^{\circ}_{Cu^{++}/Cu} = 0.34 \text{ V}$$

- a) write the cell reaction.
⇒ $E^{\circ}_{Zn/Zn^{++}} > E^{\circ}_{Cu^{++}/Cu}$. so Zn act as cathode
Cu act as anode
cell reaction ⇒ $Zn + Cu^{+2} \rightarrow Zn^{++} + Cu$

- b) construct a galvanic cell.




- c) calculate the standard emf of the cell?



⇒ standard emf $E^{\circ}_{cell} = E^{\circ}_{cathode} - E^{\circ}_{anode}$
 $= 0.34 - 0.76$
 $= -0.42$

So given reaction is non feasible.

- d) ~~why~~ can you store the $CuSO_4$ in zinc vessel?

⇒ No, because the reactivity of Zn is more than Cu so, it displaces copper in $CuSO_4$ solution,





- A → Nitro benzene
- B → Amino benzene
- C → Benzene diazonium chloride
- D → phenol
- E → 2,4,6 - Trinitrophenol


6 What are the main assumptions of CFT (Crystal field theory)? Explain, the crystal field splitting in octahedral complex.

⇒ Crystal field theory (CFT) assumes that the bond between central metal cation & ligand in a complex is purely ionic.

When negatively charged or neutral ligands having lone pair of electrons come closer to the central metal ion repulsive force created between the electrons of the d-orbitals of metal cation & the electrons of ligands.

According to C.F.T at first there is an increase in the energy of d-orbitals relative to that of the free ion just as would be the case in spherical field. The two orbitals lying along the axis get repelled more strongly than other three (d_{xy}, d_{yz}, d_{xz}). The degenerate set of d-orbitals get split into two sets: the lower energy t_{2g} set and e_g set. The energy separation is denoted by Δ_o . The actual configuration adopted is decided by the relative values of Δ_o & P . P represents the energy reqd for electron pairing in a single orbital.

- (a) If $\Delta_o < P$ then fourth electron goes to e_g .
- (b) When $\Delta_o > P$, then fourth electron pairs to t_{2g} orbital.




7 When propene reacts with HBr in presence of organic peroxide, it gives major product just opposite to Markovnikov's rule.

a) State the rule.

⇒ It states that the addition of halogen acid to unsymmetrical alkenes proceeds in such a way that the hydrogen of halogen acid adds to the double-bonded carbon having a greater number of hydrogen atoms.

b) Write a chemical eqⁿ for above statement & the major product.





c) Why is HI not given Anti-Markovnikov's addition?

⇒ Because HI bond is weak & undergoes homolysis readily to form iodine free radical. The iodine free radical has greater tendency to combine together to form iodine molecule rather than adding to the ethylenic bond.

d) What happens when major product is heated with sodium metal in presence of diether?

⇒



Q) Write down the functional isomers of carbonyl compound with molecular formula C_3H_6O . Which isomer will react faster with HCN & why?


⇒ CH_3CH_2CHO will react faster with HCN due to less steric hindrance & electronic reason than CH_3COCH_3 .

b) Alkenes ($C=C$) & carbonyl compound ($C=O$) contains a pi (π) bond but alkene show electrophilic addition whereas carbonyl compound shows nucleophilic addition reaction. Explain. (2)


⇒ Alkenes undergo electrophilic addition whereas aldehydes & ketones undergo nucleophilic addition because in alkenes the double bond joins two carbon atoms & there is no resultant polarity. While in carbonyl compounds reactions due to the polarity in the carbonyl bond that makes them vulnerable to a nucleophile, an atom that donates electrons.

G-C


9) a) How would you distinguish propan-1-ol, propan-2-ol & 2-methyl propan-2-ol by victor meyer's method?




b) What happens when phenol is treated with
i) Aqueous bromine



d) Ethanoic anhydride reacts with ammonia?



e) Ethyl ethanoate is hydrolyzed in acidic medium?



Justify with reasons.

i) Acetic acid is weaker acid than formic acid.

\Rightarrow Acetic acid has methyl group attached to carboxylic acid COOH that shows positive inductive effect & increases the electron density on the C of COOH group, thereby reduces its ~~acid~~ acidity.

ii) Amide is amphoteric in nature.

\Rightarrow Amides behave both as weak acid as well as weak base & thus show amphoteric in nature.

iii) Acyl halide is most reactive acid derivatives.

\Rightarrow Inductive electron withdrawal by Cl increases the electrophilic character of the carbonyl carbon, & increases its reactivity toward nucleophiles. Thus acyl halide is most reactive acid derivatives.

11 a) The process of determining the concentration of unknown solution is titration. Define redox titration. Calculate the molality of one litre of 93% H_2SO_4 solution (weight by volume). The density of the solution is 1.84 gm/L .

2) Those titration in which the strength of oxidizing agent is determined by using standard reducing agent or vice-versa are known as redox titration.

(Given, 93% H_2SO_4 solution (weight/volume).

It means 93 g of H_2SO_4 present in 100 ml of solution,

$$\text{weight of } H_2SO_4 = 93 \text{ g}$$

$$\text{volume of solution} = 100 \text{ ml}$$

$$\therefore \text{weight of solution} = 100 \times 1.84 = 184 \text{ g}$$

Thus,

$$\text{weight of water} = 184 - 93 = 91 \text{ g}$$

$$\therefore \text{molality} = \frac{\text{moles of } H_2SO_4}{\text{weight of water in (g)}} \times 1000$$

$$\text{moles of } H_2SO_4 = \frac{93}{98} = 0.948$$

$$\therefore \text{Molality} = \frac{0.948}{91} \times 1000$$

$$= 10.42 //$$

11/b) ⇒ Acetic acid is a weak acid & ionized as,
 $CH_3COOH \rightleftharpoons CH_3COO^- + H^+$

For weak acid,

$$[H^+] = \sqrt{K_a C}$$
$$= \sqrt{1.8 \times 10^{-5} \times 2}$$
$$= 4.2426 \times 10^{-3}$$

Now, $pH = -\log [H^+]$
 $= -\log (4.2426 \times 10^{-3}) = 2.37$

Now, pH of diluted solution,
 $pH = 2 \times 2.37 = 4.7447$

& $-\log [H^+] = 4.74$
 $[H^+] = 1.81 \times 10^{-5}$

For weak acid,

$$[H^+] = \sqrt{K_a C}$$

Squaring on both side.

$$[H^+]^2 = K_a C$$

$$C = \frac{[H^+]^2}{K_a}$$

$$= \frac{(1.8 \times 10^{-5})^2}{1.8 \times 10^{-5}}$$
$$= 1.82 \times 10^{-5}$$

Now,

$$m_1 V_1 = m_2 V_2$$

$$1 \times 1 = V_2 \times 1.82 \times 10^{-5}$$

$$\therefore V_2 = 54945.05 \text{ } \Delta$$


G-11

Date _____
Page _____

1 a) Bronsted Lowery Acid

⇒ It is define as a substance (ion or molecule) that has a tendency to donate proton.


Example,



b) Bronsted Lowery Base

⇒ It is define as a substance that has a tendency to accept proton i.e. proton donating species are acid but proton accepting species are base.

Example,



c) Lewis Acid ⇒ It is define as a substance i.e. (atom, ion or molecule) that has a tendency to accept the lone pair of electron from base through a coordinate covalent bond.

Example, FeCl_3 , Cu^{++} , Ag^+ , etc.

d) Lewis Base ⇒ A base is a species that has a tendency to donate a lone pair of electron from base through a coordinate covalent bond. Example, OH^- , CN^- , CaO , etc.

$$R = k [A]^0 [B]^2$$

$$R = k [A]^m [B]^n$$

Date _____
Page _____

e) conjugate acid & base of NH_3

\Rightarrow Ammonia or NH_3 is a base. It accepts a proton to give its conjugate NH_4^+ . Similarly, NH_4^+ loses a proton to give conjugate base NH_3 . Therefore, NH_4^+ is the conjugate acid of ammonia.

2 ~~Q~~

a) Why is blister copper called so?

\Rightarrow The slightly impure copper had impurities, mostly sulphur dioxide which bubbled up through the copper as it solidified. That gave a blistered surface and the results of the smelting were called blister of copper.

b) Write the composition of bronze.


\Rightarrow Bronze is a metal alloy that primarily contains copper and 12% tin.

c)


The prevention of rusting are:-

- i) Passing steam over red hot iron. Here a thin layer of ferrous oxide (Fe_3O_4) is deposited, which prevents the rusting. It is Bark's protection.
- ii) Oxide, phosphate or other chemical coating like Fe_3O_4 , zinc phosphate, etc.


4 a) Aldol condensation.



b) Benzoin condensation




c) Rosenmund's reaction




3. Describe the electrochemical theory for the rusting of iron. List any two methods of prevention of rusting.

⇒ The chemical corrosion of iron by oxidation with moist air to give hydrated ferric oxide is called rusting. The rusting process is accelerated by acidic oxides like CO_2 , SO_2 , NO_2 , etc.




According to the electrochemical theory, rusting of iron occurs due to formation of galvanic cell in the iron. At anode, iron atom oxidizes to Fe^{++} ion by losing 2 electrons.




The electron moves to cathode where oxygen reduces to OH^- by gaining electron.

At cathode,




The ferrous & anhydrous iron then diffuse & under the influence of dissolved oxygen, Fe^{++} is oxidized to Fe^{+++} , which then it combines with OH^- to form rust.




The prevention of rusting are:-

- i) Passing steam over red hot iron. Here a thin layer of ferrousferrous oxide (Fe_3O_4) is deposited, which prevents the rusting. It is Barf's protection.
- ii) oxide, phosphate or other chemical coatings like Fe_3O_4 , Zinc phosphate, etc.


4
a) Aldol condensation.




b) Benzoin condensation




c) Rosenmund's reaction




d) Oxidation reaction of acetaldehyde



e) Reaction of acetone with hydrazine




S Write different isomers of organic compound having formula $\text{C}_2\text{H}_7\text{N}$. How would you distinguish them by nitrous acid test?




We will get a bright coloured azo dye. But if we add nitrous acid to aliphatic amines, then effervescence of nitrogen gas is obtained. Thus, we can distinguish between ethylamine & ~~aniline~~ aniline by using azo dye test.

e) Ethyl alcohol is heated with acetic acid in presence of conc. H_2SO_4 .






6

a) Give an example of diazotisation reaction using compound A. (1)




b) How would you prepare p-hydroxyazobenzene using compound A & B.






d)



7a) Carboxylic acid contains the carbonyl group but do not show the nucleophilic addition reaction like aldehyde or ketone. Why? (2)



Due to resonance, the carbonyl group has a partial double bond character instead of the double bond. Hence they do not give the chemical properties of aldehyde & ketones (carbonyl group)

b) Arrange the following in decreasing order of their acidic strength. Explain the arrangements. 2,4,6-Trinitrophenol, p-nitrophenol, phenol, m-nitrophenol. (1+2).

8

⇒ Solⁿ Given,

~~SP. gravity =~~ $V_1 N_1 = V_2 N_2$

$V_1 N_1 = 0.03 \times 0.1 = 0.003$ equivalents

$N_1 = ?$ 10ml, volume of dil. acid neutralized.

$N_1 = \frac{0.003}{0.01} = 0.3 N$

0.3N is the normality of the 10ml of the 1L diluted solution.

mass of the solute in 10ml

$\frac{x}{49} / 0.01 = 0.3$

$x = 0.147 g$

Date: _____
Page: _____

$$0.147 \text{ g} = 10 \text{ mL}$$

$$m = 1000 \text{ mL}$$

$$m = 14.7 \text{ g}$$

$$\begin{aligned} \% C &= \frac{m}{\rho V} (100) \\ &= \frac{14.7}{1.8 \times 10 \text{ mL}} (100) \\ &= 81.67 \% \end{aligned}$$

The mass percentage for 1.8 g/mL - 86.9%.

Q 2) $\text{SP. gravity} = 1.8$

For diluted acid

$$V_1 = 10 \text{ mL}$$

$$N_1 = ?$$

$$N_2 = 30$$

$$V_2 = 0.1$$

Now, $N_1 V_1 = N_2 V_2$

$$\text{or, } 10 \times N_1 = 30 \times 0.1$$

$$\therefore N_1 = 0.3$$

For concentrated,

$$V_1 = 1000 \text{ mL}$$

$$N_1 = 0.3$$

$$V_2 = 10 \text{ mL}$$

$$N_2 = ?$$

Now, $N_1 V_1 = N_2 V_2$

$$1000 \times 0.3 = 10 \times N_2$$

$$\therefore N_2 = 30$$

Hence,

$$N = \frac{\% \times \text{SP. gravity} \times 10}{\text{Eq. wt}}$$

$$\% = \frac{30 \times 10}{10 \times 1.8}$$

$$\% = 81.67\%$$

OR

a) Is the titration redox or acid base?
=>

b) Why does $KMnO_4$ act as self-indicator?
Define indicator.

=> $KMnO_4$ solution are dark purple. When used as a titrant, as soon as the endpoint is reached & the $KMnO_4$ is in excess, the solution has a permanent pink hue. Thus $KMnO_4$ acts as its own indicator.

Indicator is a substance which is used to determine the end point in a titration.

c) $KMnO_4$ is not primary standard substance, why?
=> Because it is difficult to obtain the pure state of $KMnO_4$ as it is not free from MnO_2 . Also, the colour is so intense that it acts as its own indicator.

d) If 100cc of N/10 $KMnO_4$ solution is to be prepared in acidic condition, what mass of $KMnO_4$ is required?

e) 100ml of 0.1 $KMnO_4$

$$\frac{W}{E} = \frac{VN}{1000}$$


$$W = 0.316 \text{ gram}$$

Or, $\frac{W}{31.6} = \frac{100 \times 0.1}{1000}$ [E is $KMnO_4$ in acidic medium is 31.6]


b) A reference electrode is an electrode which has a stable & well-known electrode potential.

The standard hydrogen electrode consists of a platinum electrode coated with platinum black. The electrode is dipped in an acidic solution & pure hydrogen gas is bubbled through it.

Act as a cathode



Act as anode



10

⇒ For this reason, doctors may sometimes recommend medications called antipyretics to lower a person's temperature. Tylenol can also reduce a fever.

Penicillin are a group of antibacterial drugs that attack a wide range of bacteria.

It work as busting the cell wall of bacteria.

|| 1) Discuss cyanide process for extraction of silver from horn silver.


⇒ silver is extracted from the ore-argentite (Ag_2S). The process of extraction of silver is called as cyanide process as sodium cyanide solution is used. The ore is crushed, concentrated & heated with sodium cyanide solution. This reaction forms sodium argento cyanide.

The solution of sodium argento cyanide combines with zinc dust & forms sodium tetra cyanozincate & precipitated silver. This precipitated silver is called spongy silver.

The spongy silver is fused with potassium nitrate to obtain pure silver. Then the silver obtained is purified by electrolytic process.

b) Why does indelible ink made of silver nitrate is used to produce stain during election or polio drop campaign? Elucidate showing valid reaction.

⇒ When silver nitrate solution comes in contact of skin in presence of sunlight, it is reduced by ~~an~~ organic compound present in the skin into metallic silver producing black stain:



2a) How does the surface area of the reactant affect the rate of reaction? (1)

⇒ Surface area of the reactant alters the rate of reaction. Larger the surface area of the reactant, higher will be the rate of reaction & vice versa.

b) What is the role of temperature in the rate of reaction? (2)

⇒ The rate of chemical reaction can be changed by altering the temperature. If the temperature is increased, the reactant particles move more quickly. They have more energy.

c) What do you mean by catalyst? What is the role of catalyst in kinetics? (2)

⇒ Catalyst is a substance which can change the speed of a chemical reaction without itself undergoing any change in its mass & chemical composition.

Catalyst alters the rate of chemical reaction. Positive catalyst increase the rate of reaction but negative catalyst decreases the rate of reaction.

eventually decay by emitting a particle that transforms when they are unstable & transforms the nucleus into a lower energy state so, this ~~reaction~~ reason the α & β -rays are produced.

b) For nuclear reaction, α -particles & protons are accelerated but neutrons must be slowed down. Explain. (2)

\Rightarrow The positively charged bombarding particles called projectiles suffers repulsion therefore α & β particles should be accelerated with high energy. On the other hand, neutrons are neutral particles & these have to be slowed down so that they could be captured by the nucleus.

c) Write the medical uses of radioisotopes. (1)

\Rightarrow They can be used for imaging to study the dynamic processes taking place in various parts of the body.


5 Describe chemical method for the separation of primary, secondary & tertiary amines by Hoffmann's method. (5)

3 ⇒ The S_N1 reaction is a unimolecular substitution reaction & follows the first order kinetics i.e. the rate of reaction depends on the concentration of substrate only (Rate = k (substrate)).

The S_N2 reaction is a bimolecular substitution reaction & follows the second order kinetics i.e. the rate of reaction depends on the concentration of substrate & nucleophile.

~~As C-X bond~~

The halogen atom attached to the benzene ring in Aryl halides are slightly deactivating & also ortho-para directing. Since the ring gets deactivated as compared to that of benzene, haloarenes are less reactive than haloalkane toward nucleophilic substitution reaction.



4

a) How are α & β - rays produced?

⇒ During radioactive particles like α & β rays are emitted by an atom due to unstable atom trying to gain stability. Hence, the atoms

eventually decay by emitting a particle that transforms when they are unstable & transforms the nucleus into a lower energy state so this ~~reaction~~ reason the α & β -rays are produced.

b) For nuclear reaction, α -particles & protons are accelerated but neutrons must be slowed down. Explain. (2)

\Rightarrow The positively charged bombarding particles called projectiles suffers repulsion therefore α & β particles should be accelerated with high energy. On the other hand, neutrons are neutral particles & these have to be slowed down so that they could be captured by the nucleus.


c) Write the medical uses of radioisotopes. (1)

\Rightarrow They can be used for imaging to study the dynamic processes taking place in various parts of the body.

5 Describe chemical method for the separation of primary, secondary & tertiary amines by Hoffmann's method. (5)


OR

Identify A, B, C, D & E with their names.




6
27


8) a) calomel with ammonia.




b) corrosive sublimate with potassium thiocyanate.




c) mercuric sulphate when grinded with sodium chloride in presence of MnO_2 .



d) Mercury with excess chloride.



e) Mercury (II) chloride with potassium iodide.



OR

Give reason:

- a) Open hearth process provides better quality of steel than any other method.
- 2) Because
 - i) oxidation of impurities done by haematite ore instead of air.
 - ii) Fuel is saved by using regenerating system of heat.
 - iii) Percentage of carbon and silicon are decreased by adding scrap (unused) iron and low grade wrought iron.

b) ~~corrosive substrate with potassium thiocyanate.~~

- b) A piece of Al or Zn can partially protect iron tank from rusting.
 - ⇒ The iron or steel is coated in a thin layer of zinc. This stop oxygen and water reaching the surface metal underneath but the zinc also act as a sacrificial metal. Zinc is more reactive than iron, so it oxidises in preference to the iron object.



- c) Presence of acidic gases in the atmosphere increases the rate of rusting.
 - ⇒ Due to formation of galvanic cell.

- Date: _____
Page: _____
- d) Haematite ore is separated by gravity separation method but not with froth floatation method.
- e) Because froth floatation method is mainly used for the sulphide mineral & gravity separation method is gravity of ore and associated impurities. Hence, this oxid method is used for concentration of heavy oxide ore. Haematite ore is combination of iron & oxide & also it is heavy mineral oxide. Hence gravity separation is used.


G-C

9 ~~10~~ From alcohols:-



a) Reaction with halogen acid (HX):



b) Reaction with phosphorous halide (PX₃ or PX₅):



c) RXⁿ with thionyl chloride (SOCl₂) (Darzenski's Rⁿ):



a) What is Lucas reagent?

⇒ The mixture of conc. HCl & anhydrous $ZnCl_2$ is called Lucas reagent.

b) Is $ZnCl_2$ necessary for tertiary alcohol with HCl? If not why?

⇒ $ZnCl_2$ is not necessary for tertiary alcohol with HCl because the tertiary alcohol is more reactive.

c) Haloalkanes are slightly soluble in water. Give reason.

⇒ Due to intermolecular attraction betⁿ haloalkanes & water molecules.

d) What is the order of boiling point for n-propyl chloride, isopropyl chloride & tertiary butyl chloride? Describe with suitable reason.

⇒

Given,

b) \Rightarrow

$$E^{\circ}_{\text{Cu}^{2+}/\text{Cu}} = 0.34\text{V}$$

$$E^{\circ}_{\text{Ag}^{+}/\text{Ag}} = 0.80\text{V}$$

Now,

$$E^{\circ}_{\text{cell}} = E^{\circ}_{\text{cathode}} - E^{\circ}_{\text{anode}}$$

$$= +0.80 - 0.34$$

$$= +0.46\text{V}$$

Here,

$E^{\circ}_{\text{cell}} = +ve$. So given reaction is feasible.

The KCl - bridge cannot be used when any salt of lead, or silver is used in the cell because lead chloride & silver chloride are insoluble in water.

41
 \Rightarrow

Solution we have,

$$N = \frac{\%(\text{w/w}) \times \text{sp. gravity} \times 10}{\text{Eq. wt.}}$$

$$= \frac{24.7 \times 1.18 \times 10}{49} = 5.948\text{N}$$

From law of equivalent proportions,

No. of g. eqv. of H_2SO_4 - No. of g. eq. of aluminium
= No. of g. eqv. of remaining H_2SO_4 in dilute solution.

$$\text{or, } \frac{N_1 \times V_1}{1000} - \frac{W}{E} = \frac{N_2 \times V_2}{1000}$$

$$\text{or, } \frac{5.948 \times 75}{1000} = \frac{2.7}{27/3} = \frac{N_2 \times 400}{1000}$$

$$\text{or, } N_2 = 0.365 \text{ N}$$

or, Normality of the free H_2SO_4 in the resulting solution = 0.365 N.

Now, we have

$$\text{Normality} = \text{Molarity} \times \text{basicity}$$

$$\text{or, molarity of free } \text{H}_2\text{SO}_4 \text{ in the resulting solution} \\ = \frac{0.365}{2} = 0.182 \text{ M}$$