

## 'Nitro compound'

DATE     Introduction and Nomenclature.



Organic compounds containing nitro ( $-NO_2$ ) as the functional group are called nitro compounds. It is of two types: Aliphatic nitro compound and aromatic nitro compound

$R-NO_2$   
Aliphatic nitro compound

$R-NO_2$   
Aromatic compound

- $1^\circ$ -nitroalkanes
- $2^\circ$ -nitroalkanes
- $3^\circ$ -nitroalkanes


→ Nitro group is isomeric with Nitrite.

Nomenclature :-


→ Nitro compounds are solely (singly) named by using IUPAC system. Nitro compounds are named by using prefix nitro.

→ Accordingly aliphatic nitro compounds are called nitroalkanes while aromatic nitro compounds are called nitroarenes


→ The position of the nitro groups and other substituent are indicated by number i.e. 1, 2, 3, ... etc with the carbon atom bearing the nitro group getting the possible lowest number

CompoundsNomenclature

Nitromethane



Nitro-ethane



1-Nitro propane




2-Nitro propane




Aromatic




Nitrobenzene




2-nitrotoluene or o-nitrotoluene




1,3-dinitrobenzene.



ethyl  
2-nitrobenzene



2,4,6-trinitrotoluene (TNT)




2,4,6-trinitrophenol (picric acid)




## # Preparation of Aliphatic Nitro compounds (Nitroalkanes) (2)

1. From alkyl halides (haloalkane)  $\rightarrow$  when alkyl halides are treated with silver nitrite in presence of alcohol gives nitroalkane.




Eg:-



2. From alkanes  $\rightarrow$  This method is mainly used to prepare 1°-nitroalkanes and not applicable for prepn of aromatic nitro compounds.

2. From alkene  $\rightarrow$  when alkane is heated with fuming  $HNO_3$  in the vapour phase at 150-400°C under pressure gives nitroalkanes.

# Physical properties  $\rightarrow$ 

1. physical state  $\rightarrow$  Nitroalkanes are colorless liquids with pleasant smell.


2. Boiling points  $\rightarrow$  The nitro compounds are highly polar and thus have strong dipole-dipole interactions which cause them to have high boiling points than hydrocarbon of comparable molecular mass.

3. Solubility  $\rightarrow$  Nitroalkanes are sparingly soluble in water but soluble in organic solvents like alcohol, ether etc.

## DATE

# Chemical properties of Nitroalkanes.

**Reduction:** - Nitro compounds are reduced to primary amines through the following cases:-




**Note:** - The final product depends upon 
}


pH of reaction medium

Nature of reducing agent


a) **Reduction in acidic medium:** - when aliphatic nitro compounds are treated with active metals like Zn, Fe or Sn and conc. HCl give primary amines.

Eg:-




b) **Reduction in neutral medium** → when aliphatic nitro compounds are heated with Zn/NH<sub>4</sub>Cl (aq) gives corresponding hydroxyl amines.

Eg:-



c) **Catalytic reduction:** → when aliphatic nitro compounds are treated with H<sub>2</sub> (Ni) or H<sub>2</sub>/Pt or H<sub>2</sub>/Pd gives corresponding primary amines.


Eg:-






(d) Reduction with lithium aluminium hydride →

e.g.:-




Uses:-




[8] Aromatic - Nitro Compounds. (Nitrobenzene)

Laboratory preparation of Nitrobenzene. ( $\text{C}_6\text{H}_5\text{NO}_2$ )

When benzene is heated in the presence of conc.  $\text{HNO}_3$  and conc.  $\text{H}_2\text{SO}_4$  at  $60^\circ\text{C}$  gives nitrobenzene.



At Higher temperature further nitrated to give m-dinitrobenzene.



m-dinitrobenzene

- Procedure: At first 60 ml of conc.  $\text{HNO}_3$  & 60 ml of conc.  $\text{H}_2\text{SO}_4$  are taken in a round bottomed flask then 50 ml of benzene is added gradually to the flask, a little at a time with shaking and cooling after each addition.

After addition of all benzene the mixture is refluxed on water bath at  $60^\circ\text{C}$  for about  $1\frac{1}{2}$  hours till the yellow oily layer appears on the surface with bitter almond odour.

Purification:-

The upper layer is separated by separation funnel followed by washing with dil.  $\text{Na}_2\text{CO}_3$  to remove acidic impurities and then with water several times. After that it is dried over anhy.  $\text{CaCl}_2$  & finally distilled at about  $208^\circ\text{C} - 212^\circ\text{C}$  to obtain pure nitrobenzene.



## \* Physical properties of nitrobenzene:


1. It is pale yellow oily liquid with smell of bitter almonds.
2. It is insoluble in water but soluble in organic solvents.
3. It has bpt :  $210^{\circ}\text{C}$  and specific gravity 1.2.
4. It is highly toxic since its vapours are readily absorbed by skin.

## \* CHEMICAL PROPERTIES OF NITROBENZENE (IMPORTANT)


A. Due to nitro group:-

1. Reduction:

a) Reduction in acidic medium  $\rightarrow$  When nitrobenzene reacts with active metals like Zn, Fe or Sn in the presence of conc. HCl gives aniline.





b) Reduction in neutral medium  $\rightarrow$  when nitrobenzene reacts with  $Zn/NH_4Cl(aq)$  gives phenyl hydroxyl amine.




v. imp.

c) Reduction in alkaline medium  $\rightarrow$  when nitrobenzene reacts with  $Zn$  and  $NaOH$  gives hydrozobenzene via azoxybenzene and azobenzene.




Note:

$\rightarrow$  Thus, bimolecular reduction takes place in alkaline medium to give different products, with different reducing agent.






(d) Electrolytic reduction  $\rightarrow$  when nitrobenzene is reduced electrolytically in the presence of acid, yields phenyl hydroxylamine, which rearranges to p-aminophenol.




(e) Reduction with lithium aluminium hydride  $\rightarrow$  when nitrobenzene reduced with  $\text{LiAlH}_4$ , yields azobenzene.



[B] Electrophilic substitution reactions  $\rightarrow$  The nitro group is electron withdrawing group and therefore causes electron deficiency at ortho and para position due to -I effect and it is found that the  $e^-$  density is relatively greater in meta position because no positive charge develops on it. therefore, electrophile attacks in meta position.

Hence nitro group is meta directing group or meta director.

$\rightarrow$  the nitro group is ring deactivating so rxn takes place in drastic condition.




Uses:-

- 1) For manufacture of aniline.
- 2) For various aromatic compounds used in detergents, dyes and other pharmaceutical.
- 3) Manufacture of explosive eg:- TNT, TNBC 1,3,5-Trinitrobenzene ROX (Research and Development explosives)
- 4) Used as high boiling inert solvent.
- 5) Used as scent for cheap soaps.


Question

Conversion:-


1. Nitrobenzene to phenol:-




2. Nitrobenzene to p-Nitroaniline:-



3. Nitrobenzene to Anisole:->




classmate






4. Nitrobenzene to P-aminobenzene.



5. Toluene to 1,3,5-trinitrobenzene.



1,3,5-Trinitrobenzene.







# Bipin Khatri


(Bipo)

---

**Class 12** complete notes and paper collection.

Folders Name ↑

 Biology	 chemistry
 English	 maths
 Nepali	 Physics

 Drive

---

Feedbacks:

[admin@bipinkhatri.com.np](mailto:admin@bipinkhatri.com.np) | [bipinkhatri.ram@gmail.com](mailto:bipinkhatri.ram@gmail.com)

---

Contact:



@im.bipo



[www.bipinkhatri.com.np](http://www.bipinkhatri.com.np)



@im.bipo